


Investigating differences between cirrus optical depth retrievals between CALIOP V3 and MODIS observations

Robert Holz, Steve Platnick, Ralph Kuehn,
and Mark Vaughan





What Was Compared

- 4 months of collocated daytime CALIOP and MODIS cirrus optical depths (MODIS C5 and CALIOP V3)
- Single layer, CALIOP non-opaque
- Non-Polar ocean (<60 deg)
- Phase determined by CALIOP V3 phase retrieval






MODIS vs CALIOP OD






MODIS vs CALIOP OD






IR Cloud Forcing Difference


Feb 2007







IR Radiative Closure

- Model: LBRTM/DISORT (LBLDIS)
- Ping et al. scattering calculations
- Surface/Thermodynamic profile: GDAS
- Cloud boundaries and phase: CALIOP
- Cloud effective radius: MODIS
- **Variable:** MODIS/CALIOP layer OD




IR Closure Unconstrained





Bias Hypothesis

- MODIS
 - Uncertainties in the single-scatter calculations used in the OD retrieval (asymmetry parameter)
- CALIOP
 - Multiple scattering
- Non-uniform cloud optical properties




MODIS Scattering Properties


- To first order the MODIS retrieval of optical depth is:

$$R_v = (1 - g) \times \tau$$



- For ice, g is determined by relating MODIS spectral observations to theoretical single scatter calculations
- **Question:** Are these calculations representative of real ice crystals?


0.86 μm MODIS C5 g vs. r_e








MODIS Retrieval Space





CALIOP Multiple Scattering





Conclusions

- A factor of two difference was found between MODIS and CALIOP retrieved OD for thin cirrus ($OD < 3$)
- IR radiative closure study concluded neither MODIS or CALIOP OD agree with measured TOA MODIS 11 μm BT
- Possible sources of uncertainty have been identified in both CALIOP and MODIS
- A community effort is necessary to understand and resolve the OD differences